



STUDY OF PHYSICAL FACTORS AFFECTING THE DENSITY OF SEA SNAILS. HAT SAMRAN BEACH TRANG, THAILAND

Researchers Team: Miss Warapha Bamrung, Miss Chanisa Aunkaew, Miss Nichapat Sriphon Grade Level : 11th Grade

Advisor: Teacher Sawitree Duangsook Wichienmatu School, Trang Province



Abstract

The study of physical factors affecting the density of wedge clams on Samran Beach, Hat Samran District, Trang Province aimed to investigate the physical factors influencing the density of wedge clams in the area of Samran Beach, Hat Samran District, Trang Province. The study examined the density of wedge clams, water temperature, pH, dissolved oxygen, and salinity to understand the density of wedge clams in the Samran Beach area, Hat Samran District, Trang Province. The results showed that the density of wedge clams on Samran Beach, Hat Samran District, Trang Province was influenced by physical and chemical factors. The water temperature ranged from 29.4-33.3 °C, which is conducive to the survival of aquatic organisms. The pH ranged from 5.6-6.8, considered normal and suitable for aquatic life. The dissolved oxygen (DO) ranged from 4.29-7 mg/l, and the salinity ranged from 28.4-30.64 ppt. The cloud cover showed that Stratocumulus clouds were more prevalent than Cirrus clouds, indicating high humidity or changing weather conditions.

Keywords: Wedge clams, Samran Beach, Hat Samran District, Trang Province, Physical factors, Wedge clam density

Research Questions

Are the physical factors affecting the density of wedge clams in the Samran Beach area, Hat Samran District, Trang Province the same?

Introduction

Background and Significance of the Project

The wedge clam is a bivalve mollusk belonging to the order Cardiida, family Donacidae. In Thailand's coastal areas, there are 8 confirmed species (Signorelli and Printrakoon, 2019). They have a thick, triangular shell with the upper edge of each shell longer on the front side than the back. The lower edge of the shell is circular, and the apex points towards the rear of the body. The outer shell surface has relatively smooth horizontal lines and comes in various colors such as white, cream, brown, reddish-brown, clear white, light yellow, black, or purplish-brown. These animals live in the intertidal zone on sandy beaches (Swennen et al., 2001), an area subject to constant temperature changes, allowing them to adapt well to both tropical and temperate climates. Wedge clams feed on plankton and organic matter suspended in seawater. The lower Andaman Sea coastline features long sandy beaches stretching north-south from Ranong, Phang Nga, Phuket, Trang, and Satun provinces, an area rich in marine resources and important for biodiversity, filtering food from seawater. Samran Beach is located in the lower Andaman region, Moo 9, Phum Ban Na-Hat Samran Road, Hat Samran Subdistrict, Hat Samran District, Trang Province 92120. From Samran Beach, you can see Libong Island, Lao Liang Nuea Island, Lao Liang Tai Island, Petra Island, and Sukorn Island. It boasts black sand beaches, numerous minerals, and mud suitable for spa treatments, making it a tourist destination with abundant natural resources. It is also significant in terms of biodiversity (Department of Marine and Coastal Resources, 2022). Therefore, the researchers are interested in studying the physical factors affecting the density of wedge clams at Samran Beach, Hat Samran District, Trang Province

Research Methods

1. Research Preparation

- 1) Define the research topic.
- 2) Conduct literature review and gather relevant knowledge and theories.
- 3) Define the research objectives.
- 4) Determine the sampling points within the study area.

2. Processing floor

- 1) Conduct research operations planning.
- 2) Conduct a survey of the area to be researched.
- 3) Collect soil and water samples to measure soil and water temperature using a thermometer. Read the values and record the results.
- 4) Measure the salinity of water with a Salinity Meter, read the words and record the results.
- 5) Measure the acidity-alkalinity (pH) of the soil with a universal indicator, read the value and record the results.
- 6) Analyze the values obtained from the survey and summarize the study results.


Results

Geographic coordinates: Study of the area around Samran Beach, Hat Samran Subdistrict

Hat Samran District, Trang Province 92120 has coordinates 7.2352 °N, 99.5366 °E.

Table 1 Geographic coordinates

สถานที่เก็บตัวอย่าง	พิกัดทางภูมิศาสตร์	
	Latitude (N)	Longitude (E)
ชายหาดสำราญ	7.2352	99.5366



Study of the physical quality of water

Table 2 Water temperature

Location	Trial 1	Trial 2	Trial 3	Average (X) ± S.D
1	28.35	29.4	30.01	29.25 ± 0.84
2	29.45	29.3	30.5	29.75 ± 0.65

From Chart 2 showing the water temperature, it was found that the average temperature of area 1 was 29.25±0.84 and the average temperature of area 2 was 29.75±0.65.

Chemical study of water quality

Table 3 Water salinity, pH, and dissolved oxygen content.

Property	Location 1	Location 2	Location 3	Location 4	Location 5	Average (X) ± S.D
1. Salinity (ppt)	28.2	28.6	28.4	28.2	28.6	28.40 ± 0.2
2. pH	7	7	7	7	7	6.80 ± 0.05
3. Dissolved Oxygen (mg/L)	7.5	7.5	7	7	6	7.00 ± 0.61

From the data in Table 3, the salinity of water, the acidity (pH), the amount of oxygen dissolved in water, the average seawater saturation value was 28.4±0.2 ppt, the average pH value was 6.8±0.45, the amount of oxygen (DO) was 7.00±0.61 mg/L

Soil quality study

Table 4 Soil texture and color

Type	Location 1	Location 2	Location 3	Location 4	Location 5
Texture	Sandy clay loam	Sandy clay loam	Clay	Sandy loam	Clay
Color	Dark brown	Dark brown	Dark brown	Dark brown	Dark brown

From Table 4, soil quality found that in the area with sea shells, there are both clay and sandy soil, sandy soil, loam soil and clay soil, and all areas of soil are dark brown.

Study of the density of sea snails

Table 5. Density of sea clams.

Type	Location 1	Location 2	Location 3	Location 4	Location 5	Average (X) ± S.D
Number (clams)	22	23	10	18	6	15.80 ± 7.50
Density (clams/cm²)	0.61	0.63	0.27	0.5	0.16	0.43 ± 0.20

From Table 5, the density of all sea clams in Samran Beach area showed that the study area had a density of sea clams of 0.43±0.20 individuals/square meter.

Table 6 Analysis of cloud cover in areas with and without shellfish.

Type	Trial 1	Trial 2	Average (X) ± S.D
Stratocumulus	72.50%	12.50%	42.50 ± 42.42
Cirrus	10%	35%	22.50 ± 17.67

From Table 6, showing the amount of cloud cover, it was found that Stratocumulus cloud cover was more than Cirrus cloud cover. Which indicates high altitude weather or changes in weather conditions.

Discussion

From the study of physical factors affecting the density of sea clams in the area of Samran Beach, Hat Samran District, Trang Province, it was found that the important factors affecting the density of sea clams are the physical characteristics of the beach surface. Sea clams are mostly found in areas with clay mixed with sand and salinity values between 28.2-28.6 ppt, pH = 7 and dissolved oxygen value of 7.5 mg/l, which affects the density of sea clams. A significant change in salinity may cause the number of sea clams to decrease. Temperature and environmental factors. The sea temperature that is suitable for sea clams to live is between 26-30 degrees Celsius. Changes in weather conditions, such as heavy rain or strong winds, affect the density of sea clams. The rate of sea water rise and fall (ebb and flow) Sea clams are distributed along the beach during low tide and are found to be denser in areas that receive continuous seawater.

Conclusion

Therefore, the factors affecting the density of sea clams, including temperature, salinity, pH and dissolved oxygen, affect the density of sea clams. This study helps to understand the ecosystem of Samran Beach and can be applied to manage sea clam resources for conservation and sustainable utilization.

Bibliography

Morphological variation of the clam, KKU Science, 2021.

<https://ph01.tci-thaiho.org/index.php/KKUSCIJ/article/download/250287/169753/900172>

Some Ecology of the Snail (2018)

<https://scijournal.buu.ac.th/index.php/sci/article/view/2571/2276>

Information about snails

<https://th.m.wikipedia.org/wiki/%E0%B8%AB%E0%B8%AD%E0%B8%A2%E0%B9%80%E0%B8%AA%E0%B8%B5%E0%B8%A2%E0%B8%9A>

Some ecology of clams

Some ecology of clams

<https://scijournal.buu.ac.th/index.php/sci/article/view/2571/2276>

Diversity and density of sea shells in Ao Boonkhong, Sikao District, Trang Province

<https://search.app/ByivndTillIpyaVE09>