



Title of Research

A Study of Air Temperature and Cloud Types at Ban Thung Na School, Trang Province

Ban Thung Na School (Mueang)



Abstract

This environmental science research aimed to: (1) compare air temperatures at different times of the day at Ban Thung Na School, Trang Province, and (2) compare cloud types observed at different times of the day at the same location. Air temperature measurements and cloud observations were conducted at three different times. The results showed that the highest air temperature occurred at 12:00 p.m., and the greatest variety of cloud types observed was four types: Stratus, Cumulus, Cirrus, and Altocumulus.

Keywords: Air temperature, Cloud types

Research Question

1. Does air temperature differ at different times of the day? How?
2. Do cloud types differ at different times of the day? How?

Introduction

Clouds are diverse and can be classified by altitude, shape, and thickness. Clouds directly affect air temperature by acting as a protective layer and helping maintain the Earth's heat balance. Thick white clouds reflect sunlight, causing the Earth's surface to cool, while some thin clouds help trap heat beneath them. Cloud formation from water vapor condensation releases latent heat, and cloud dissipation causes the temperature within the cloud to equilibrate with the surrounding atmosphere. Therefore, this study focuses on air temperature and cloud types at Ban Thung Na School, Trang Province.



Field Photos
(requires release forms)

Research Hypotheses

1. Air temperature differs at different times of the day.
2. Cloud types differ at different times of the day.

Study Area

The study area was Ban Thung Na School, Mueang District, Trang Province, located at latitude 7.6936 and longitude 99.7388. The area is characterized by atmospheric conditions and cloud cover over the school. Air temperature and cloud types were measured and observed at this location.

Data Collection

Data were collected following standard air measurement protocols related to air temperature and cloud types at different times.

Data Collection Period:
November to December 2025



Materials and Equipment

1. Cloud chart
2. Hygrometer
3. GLOBE Observer App

Methodology

Part 1: Study of Air Temperature at Different Times

1. A thermometer was installed in a Stevenson screen and reset, then left in place for one day.

2. Temperature readings were taken every 15 days for a total of three sessions. In each session, temperatures were recorded at three times: 9:00 a.m., 12:00 p.m., and 3:00 p.m. Observations were recorded and results documented.

Part 2: Study of Cloud Types at Different Times

Cloud charts were studied and used to observe cloud types in the sky every 15 days for a total of three sessions. In each session, cloud photographs were taken at three times: 9:00 a.m., 12:00 p.m., and 3:00 p.m., and the results were recorded.

Conclusion and Discussion

The environmental science study on air temperature and cloud types at Ban Thung Na School, Trang Province, found that the highest air temperature occurred at 12:00 p.m. The greatest variety of cloud types observed was four types: Stratus, Cumulus, Cirrus, and Altocumulus.

Acknowledgements

This environmental science research on air temperature and cloud types at Ban Thung Na School, Trang Province, was successfully completed with the support and encouragement of the school director, Mr. Thirasanti Kanchanasrimek. The researchers would like to express their sincere gratitude to Mrs. Kwanjai Kanchanasrimek for her kindness, assistance, guidance,